

—

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] The subject application is a Continuation of application No. 08/450,962, filed May 25, 1995, now Pat. No. 6,274,706, which is a Division of application No. 08/117,829, filed Sept. 08, 1993, now abandoned, which is a Continuation-in-Part of application No. 07/721,571, filed on Sept. 03, 1991, now Pat. No. 5,288,607.

BACKGROUND OF THE INVENTION

[0002] The present invention relates to the entire gene encoding beta ("β") 3-adrenergic receptor polypeptides of mouse and human origin, which polypeptides provide a procedure for studying the effects of various chemical agents on the β3-adrenergic receptor and coupled adenylate cyclase and hormone-sensitive lipases. More particularly, the present invention also relates to antibodies, vectors, nucleotide probes, cell hosts transformed by genes encoding polypeptides, having β3-adrenergic receptor activity.

[0003] In the past, two main classes of adrenergic receptors have been identified as the α adrenergic receptors and the β adrenergic receptors. These adrenergic receptors produce various responses to effector organs such as the eye, heart, arteriols, veins, lungs, stomach, intestine, gallbladder, kidney, skin, spleen, liver and pancreas to mention a few. These specific receptors have been defined by the effects of particular synthetic agonists which stimulate the receptors biological function and antagonists which block the adrenergic receptors biological function.

[0004] Within the two classes of adrenergic receptors, four subtypes, α_1 , α_2 , β_1 and β_2 , of these receptors for catecholamines have been identified. See, Cotecchia et al., P.N.A.S., 85, pgs. 7159-7163 (1988); Kobilka et al., Science 238, pgs. 650-656 (1987); Frielle et al., P.N.A.S., 84, pgs. 7920-7924 (1987); and Emorine et al., P.N.A.S., 84, pgs. 6995-6999 (1987). Drugs that selectively block or stimulate one of these receptor subtypes are used

096611
096612
096613
096614
096615
096616
096617
096618
096619
096620
096621
096622
096623
096624
096625
096626
096627
096628
096629
096630
096631
096632
096633
096634
096635
096636
096637
096638
096639
096640
096641
096642
096643
096644
096645
096646
096647
096648
096649
096650
096651
096652
096653
096654
096655
096656
096657
096658
096659
096660
096661
096662
096663
096664
096665
096666
096667
096668
096669
096670
096671
096672
096673
096674
096675
096676
096677
096678
096679
096680
096681
096682
096683
096684
096685
096686
096687
096688
096689
096690
096691
096692
096693
096694
096695
096696
096697
096698
096699
096700
096701
096702
096703
096704
096705
096706
096707
096708
096709
096710
096711
096712
096713
096714
096715
096716
096717
096718
096719
096720
096721
096722
096723
096724
096725
096726
096727
096728
096729
096730
096731
096732
096733
096734
096735
096736
096737
096738
096739
096740
096741
096742
096743
096744
096745
096746
096747
096748
096749
096750
096751
096752
096753
096754
096755
096756
096757
096758
096759
096760
096761
096762
096763
096764
096765
096766
096767
096768
096769
096770
096771
096772
096773
096774
096775
096776
096777
096778
096779
096780
096781
096782
096783
096784
096785
096786
096787
096788
096789
096790
096791
096792
096793
096794
096795
096796
096797
096798
096799
096800
096801
096802
096803
096804
096805
096806
096807
096808
096809
096810
096811
096812
096813
096814
096815
096816
096817
096818
096819
096820
096821
096822
096823
096824
096825
096826
096827
096828
096829
096830
096831
096832
096833
096834
096835
096836
096837
096838
096839
096840
096841
096842
096843
096844
096845
096846
096847
096848
096849
096850
096851
096852
096853
096854
096855
096856
096857
096858
096859
096860
096861
096862
096863
096864
096865
096866
096867
096868
096869
096870
096871
096872
096873
096874
096875
096876
096877
096878
096879
096880
096881
096882
096883
096884
096885
096886
096887
096888
096889
096890
096891
096892
096893
096894
096895
096896
096897
096898
096899
096900
096901
096902
096903
096904
096905
096906
096907
096908
096909
096910
096911
096912
096913
096914
096915
096916
096917
096918
096919
096920
096921
096922
096923
096924
096925
096926
096927
096928
096929
096930
096931
096932
096933
096934
096935
096936
096937
096938
096939
096940
096941
096942
096943
096944
096945
096946
096947
096948
096949
096950
096951
096952
096953
096954
096955
096956
096957
096958
096959
096960
096961
096962
096963
096964
096965
096966
096967
096968
096969
096970
096971
096972
096973
096974
096975
096976
096977
096978
096979
096980
096981
096982
096983
096984
096985
096986
096987
096988
096989
096990
096991
096992
096993
096994
096995
096996
096997
096998
096999
097000
097001
097002
097003
097004
097005
097006
097007
097008
097009
097010
097011
097012
097013
097014
097015
097016
097017
097018
097019
097020
097021
097022
097023
097024
097025
097026
097027
097028
097029
097030
097031
097032
097033
097034
097035
097036
097037
097038
097039
097040
097041
097042
097043
097044
097045
097046
097047
097048
097049
097050
097051
097052
097053
097054
097055
097056
097057
097058
097059
097060
097061
097062
097063
097064
097065
097066
097067
097068
097069
097070
097071
097072
097073
097074
097075
097076
097077
097078
097079
097080
097081
097082
097083
097084
097085
097086
097087
097088
097089
097090
097091
097092
097093
097094
097095
097096
097097
097098
097099
097100
097101
097102
097103
097104
097105
097106
097107
097108
097109
097110
097111
097112
097113
097114
097115
097116
097117
097118
097119
097120
097121
097122

extensively in clinical medicine. Despite the efficacy of these drugs, many produce side effects in individuals, due to their interaction with other receptor subtypes. For identification of the various drugs which act on the receptor subtypes, see Goodman and Gilman's The Pharmacological Basis of Therapeutics, 8th edition, MacMillan Publishing Co., Inc. (1990).

[0005] The analysis of the genes of these receptors indicate that all of the subtypes of the receptors belong to the family of integral membrane receptors exhibiting striking homologies, in particular these homologies are present in the 7 transmembrane regions. These receptors are coupled to regulatory proteins termed G proteins which are capable of binding molecules of guanosine triphosphate (GTP).

[0006] The G proteins have the capacity to intervene structurally and functionally between receptors and enzymes catalyzing the production of intracellular mediators such as adenylate cyclase, guanylate cyclase, the phospholipases and the kinases, or between receptors and ion channels. Thus, the G proteins have the capacity to regulate the flux of ions such as calcium, potassium, sodium and hydrogen ions.

[0007] The above-mentioned subtypes of receptors are known in the art as the "R₇ G family" as described by Emorine et al. Proc. NATO Adv. Res. Workshop (1988). The R₇ family comprises acetylcholine muscarinic receptors, serotonin receptors, the receptors for neuropeptides, substance K, angiotensin II and the visual receptors for the opsins, as described by Dixon et al Annual Reports in Medicinal Chemistry, pgs. 221-233 Ed. Seamon, K. B., Food and Drug Administration, Bethesda, Md. (1988) and Emorine et al., supra.

[0008] Recently, a third subtype of the β -adrenergic receptor termed β 3-adrenergic receptor, has been identified and characterized in humans and in rodents, which polypeptide does not have a receptor activity similar to that of the β 1- or β 2-adrenergic receptors. This "new" β 3-receptor in humans has been identified and sequenced as described in French application No.8900918 filed Jan. 25, 1989 and PCT/FR90/00054, resulting in U.S. application Ser. No.07/721,571, filed Jan. 25, 1990, U.S. Pat. No. 5,288,607, incorporated herein by reference, as containing 402 amino acids, which is capable of activating adenylate cyclase in the presence of an agonist, which activity increases in the order of agonists of salbutamol, BRL 28410, BRL 37344 and (1)-isoproterenol. Antibodies directed to the

100-443886-100

[0010] A pharmacological comparison of the mouse with the human counter part of the β 3-adrenergic receptors indicates that these receptors have similar reactivity. For example, mouse β 3-adrenergic receptor can be activated by CGP 12177A, oxprenolol and pindolol while displaying a low stereoselectivity and characteristic potency order for full agonists, which is similar to the human.

[0012] Thermogenesis and lipolysis in brown and white adipose tissues are under the control of this β 3-adrenergic receptor subtype as described by Arch et al., Proc. Nutr. Sci., 48, pgs. 215-223 (1989); Arch et al., In Obesity, John Wiley & Sons Ltd., London (1991); and Zaagsma et al., Trends Pharmacol. Sci., 11, pgs. 3-7 (1990). Besides adipose tissues, the expression of the β 3-adrenergic receptor has also been reported in various other tissues such as tissues of the digestive tract, from oesophagus to colon and in the gallbladder. See, for example, Bond et al., Br. J. Pharmacol., 95, pgs. 723-734 (1988); Coleman et al., British Journal of Pharmacology Proc. Supl., 90, 40 (1987); Bianchetti et al., Br. J. Pharmacol., 100, pgs. 831-839 (1990); Granneman et al., J. Pharmacol. Exp. Ther., 256, pgs. 421-425 (1991);

and Krief et al., J. Clin. Invest., (1993) (in press). This β 3-adrenergic receptor subtype has been suggested to participate in the control by catecholamines of body energy balance from intestine assimilation to storage and mobilization in adipose tissue.

[0013] Factors such as temperature, feeding or fasting and stress influence the body's hormonal status, thus inducing tissue specific adaptive modifications of energy balance which may in part result from regulation of cellular β 3-adrenergic receptor sensitivity. It has been further shown that β -adrenergic agonists, glucocorticoids and several other agents can modulate β 3-adrenergic receptor density, responsiveness and mRNA levels. The molecular determinants responsible for the regulation of cellular β 3-adrenergic receptor may possibly be found either on the receptor itself or in its gene or mRNA. For example, post-translational modifications of the receptor in the third intra-cytoplasmic loop or in the carboxy-terminal tail may modulate β -adrenergic receptor coupling to adenylate cyclase. Factors which act on β -adrenergic receptor gene transcription rate or on mRNA stability may also modulate cellular adrenergic responsiveness by modifying β -adrenergic receptor expression levels.

[0014] The β 3-adrenergic receptors have been sequenced and identified, as well as compared pharmacologically with other known receptors in mouse and humans. A comparison of the β 3-adrenergic receptor amino acid sequences predicted from the nucleotide sequences of the human and mouse genomic genes revealed differences in the carboxy-terminal regions of the receptors. Although these differences could be attributed to evolutionary species-related variations, it was recently discovered by the present inventors that the entire coding sequence for the previously reported β 3-adrenergic receptor in mouse and human was not complete. The genes encoding β 1- and β 2- comprise one exon and it was thought that the β 3-adrenergic receptor gene, when initially identified, also comprised only one exon. However, it was recently and unexpectedly discovered that contrary to β 1- and β 2-adrenergic receptor genes, the human and mouse β 3-adrenergic receptor genes comprise several exons. The identification of the introns and exons in mouse and human β 3-adrenergic receptors and thus the entire gene itself, permits a full characterization of this receptor which will aid in more sensitive genetically engineered products such as nucleotide probes, polyclonal and monoclonal antibodies and the like for drug testing and diagnostic purposes,

and permits the understanding of β 3-adrenergic receptor regulated expression in various tissues.

SUMMARY OF THE PRESENT INVENTION

[0015] Thus, it is an object of the present invention to provide polypeptides having β 3-adrenergic receptor activity in mammals which polypeptides encode the entire human and mouse β 3-adrenergic receptor genes.

[0016] A specific object of the present invention is to provide the entire amino acid and nucleotide sequences, as well as the intron/exon organization for the human and mouse β 3-adrenergic receptor which molecule characterizations are useful in not only detecting the β 3-adrenergic receptor in different species, but also testing new drugs for regulating β 3-adrenergic receptor activities.

[0017] Yet another object of the present invention is to provide nucleotide probes capable of hybridizing to the genes encoding the polypeptides having β 3-adrenergic activity in mammals for detecting this receptor in different mammalian species and for measuring the variations in the levels of expression of the β 3-receptor in mammalian cells.

[0018] Another object of the present invention is to provide antibodies, polyclonals and monoclonals which recognize the entire sequence of the β 3-adrenergic receptor in mammals, which antibodies do not recognize the β 1-adrenergic receptor or β 2-adrenergic receptor for detecting and diagnostic purposes.

[0019] A further object of the present invention is to provide recombinant vectors for the cloning and expression of β 3-adrenergic receptor proteins in mammals.

[0020] Another object of the present invention is to provide a cell host which comprises the elements of regulation making possible the expression of the nucleotide sequence encoding the polypeptides of β 3-adrenergic receptors in mammals.

BRIEF DESCRIPTION OF THE DRAWINGS

[0021] FIG. 1 depicts the nucleotide sequence (SEQ ID NO:1), amino acid translation (SEQ ID NO:2) and intron/exon organization of the human β 3-adrenergic receptor gene.

[0022] FIG. 2 depicts the nucleotide sequence (SEQ ID NO:3), amino acid translation (SEQ ID NO:4) and intron/exon organization of the mouse β 3-adrenergic receptor gene.

[0023] FIG. 3 is the complete amino acid sequence (SEQ ID NO:5) for the human β 3-adrenergic receptor.

[0024] FIG. 4 is the complete amino acid sequence (SEQ ID NO:6) for the mouse β 3-adrenergic receptor.

[0025] FIG. 5 is a schematic representation of human and mouse β -adrenergic receptor mRNA splicing.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0026] In particular, the present invention provides various genetically engineered products such as nucleotide probes, monoclonal and polyclonal antibodies for testing a variety of chemical agents including drugs, ligands and the like which may influence the regulation of β 3-adrenergic receptors. Moreover, in this testing method, it is of importance to have the entire sequence of the β 3-receptor gene, since elements in the promoter region, in the intron/exon region and in the 3' untranslated region may influence the expression of β 3. By testing drugs that may increase and/or decrease the expression levels of the β 3-adrenergic receptor, various diseases such as obesity, diabetes and hyperlipidemia and the like, may be controlled.

phosphoramidite method described in Bioorganic Chemistry, 4, pgs. 274-325 (1986) which method is suitable for the preparation of a nucleotide sequence containing a maximum of 200 nucleotides and those described in P.N.A.S., 80, pgs. 7461-7465 (1983) for nucleotides longer than 200.

[0034] Besides synthesis of nucleotides, the present invention also relates to the purification of nucleotides from genomic DNA or cellular RNA from any mammalian tissue expressing the β 3-adrenergic receptor including adipose, muscular, hepatic, intestinal, gallbladder tissues and the like. mRNA can then be isolated and cDNA synthesized therefrom by the methods described in Sambrook et al. Molecular Cloning, supra.

[0035] The nucleotide sequences or fragments thereof can be cloned and/or expressed in a plasmid, cosmid or phage type vector. Thus, the present invention also includes recombinant vectors which which can be use to clone or express in a variety of host microorganisms the β 3-adrenergic receptor. Various vectors include, but are not limited to pBr322, pUC18/pUC19, pUC119, p5P64/p5P65, pGEM-3/pGEM-4, pGEM-3Z, π AN13, Bluescript M13, λ gt10, λ 2001, λ DASH, λ FIX, C2RB, pWE15 and the like.

[0036] A preferred vector includes a HindII-PstI fragment of genomic DNA derived from mouse having about 5.5 kilobases and containing the totality of the gene coding for the β 3-adrenergic receptor in mouse including the promoter region and Exon 1, Exon 2 and Exon 3, which fragment was inserted into a HindII-PstI site of pUC18. The plasmid was transformed in E. coli(JM101). This vector, (030 B315-I) was deposited under the No. I-1272 on Dec. 1, 1992 with the C.N.C.M., Collection Nationale de Cultures de Microorganismes, 25 rue du Docteur Roux, Paris France.

[0037] A second preferred vector includes a BglII fragment of genomic DNA derived from humans having about 4.5 kilobases and containing the totality of the gene coding for the β 3-adrenergic receptor in humans including the promoter region and Exon 1 and Exon 2, which fragment was inserted into a BamHI site of pUC18. The plasmid was then transformed in E. coli (JM101). This vector, (119 B315-III) was deposited under the No. I-1273 on Dec. 1, 1992 with the C.N.C.M., Collection Nationale de Cultures de Microorganismes, 25 rue du Docteur Roux, Paris France.

[0038] For expression the vector contains in at least one of its sites not essential for replication, elements necessary to promote the expression of the polypeptide of the present invention in a cell host. For example, a promoter recognized by the RNA polymerase of the cell host may be used. Any promoter that accomplishes expression can be used, but it is preferable to use an inducible promoter and if necessary, a signal and an anchoring sequences. For expression in eukaryotic cells, the regulatory elements may include the endogenous promoter for the adrenergic receptors or viral promoters such as those of SV40 virus or Rous Sarcoma virus, while for bacterial expression, the regulatory elements may include the lactose operon, the tryptophan operon and the like.

[0039] Cell hosts transformed by any of the above-described vectors are also encompassed by the present invention. Any cell host capable of expressing any mammalian β 3-adrenergic receptor polypeptide or fragments thereof is encompassed by the present invention. These cell hosts include any eukaryotic or prokaryotic cells including bacteria such as E. coli, yeast, CHO cells and the like. It is of interest to note that when the entire intron/exon sequence of the mouse or human β 3-adrenergic receptor is expressed in a variety of eukaryotic cell hosts including CHO and L-cells, splicing of the gene takes place correctly.

[0040] The recombinant vectors and cell hosts transformed with these vectors are useful tools to generate with ease and facility various nucleotide probes.

[0041] These nucleotide probes can also be prepared from the nucleotide sequences, as discussed above, via synthesis and purification techniques and then labelled with a detectable marker. Any detectable marker can be used to label the specific nucleotide sequence of interest, such as ^{14}C , ^{32}P , enzyme markers and the like. The labelling conditions for the sequences are well known in the art and for reference such techniques described by Sambrook et al. Molecular Cloning, supra may be used.

[0042] More particularly, it is advantageous to use a full length probe having the nucleotide sequence as defined in FIGS. 1 (SEQ ID NO:1) and 2 (SEQ ID NO:3) to probe a genomic or cDNA library of different mammalian species to obtain the related β 3-adrenergic receptor of interest. A fragment of the nucleotide probe can also be generated.

[0043] A nucleotide probe is also useful in measuring the variations in the levels of expression of the β 3-adrenergic receptor in mammalian cells, since altered levels of expression may result in specific abnormalities in various mammalian tissues. More specifically, the nucleotide probes may be produced such that they react with specific sequences of the β 3-adrenergic receptor gene. Thus, even if this receptor is present in the specific species of mammals being tested, the ligand binding sites or other functional regions in the sequence may contain an abnormality that can be ascertained by the use of such probes.

[0044] The nucleotide probes which are suitable for use in the present invention include DNA probes, RNA probes, synthetic oligonucleotide probes, thioester probes and the like. Tandem probes may also be used. The tandem probes may be constructed by covalently joining two or more type-specific β 3-adrenergic receptor nucleotide sequences within a single vector. A mixture of the above-mentioned probes is also contemplated for detection purposes.

[0045] The conditions required for hybridization and washing of the oligonucleotide probes for detecting the presence of the β 3-adrenergic receptor may be different depending on the type of probe utilized. Empirically determined formulae available in the literature allow for the estimation of the oligonucleotide dissociation temperature. Hybridization of oligonucleotides also depend on several factors, which include the length of the probe and the GC content. The hybridization conditions, thus will be adjusted according to the probes used and are well within a person skilled in the art's knowledge.

[0046] The labelled hybrids are then detected depending on the type of labelled probe used. For example, an enzymatically labelled probe will require enzymatic detection, while if a radioactive label is used, gamma or beta counters or autoradiography may be used.

[0047] The present invention also relates to antibodies directed against the polypeptides of the β 3-adrenergic receptor in mammals. These antibodies are specific for the β 3-adrenergic receptor and not the β 1- nor β 2-adrenergic receptors. These antibodies may be produced by injecting the β 3-adrenergic receptor protein or polypeptide thereof into an animal. The antibodies generated by the immune response may be recovered and purified by methods known in the art. Besides their use for detection and in vitro diagnostic purposes, the

antibodies of the present invention may also be used to purify related receptors in semi-purified form by their attachment to a solid support such as gel beads.

[0048] Monoclonal antibodies may be produced by the known method described by Kohler and Milstein, Nature, 256, page 495 (1975). These antibodies are useful to diagnose the amount and the presence of the β 3-adrenergic receptor in a variety of mammalian species. Thus, the quantity and presence of this receptor may be important to diagnose diseases related to energy abnormalities in a variety of mammalian tissues.

[0049] Although direct detection of the β 3-adrenergic receptor polypeptide is possible by the use of genetically engineered products as described above, these products are not limited in use to only detecting the presence of β 3-receptors in various species. Testing of different drugs for the treatment of obesity, fatty diabetes, hyperlipidemias and the like are within the scope of the present invention by use of the aforementioned nucleotide probes, β 3-adrenergic receptors and anti-receptor antibodies.

[0050] A procedure for studying the affinity of a β 3-adrenergic receptor polypeptide for one or more chemical agents such as drugs, ligands and the like is also encompassed by the present invention.

[0051] To study the effects of various chemical agents, a cultured transformed host cell having a sequence encoding the entire β 3-adrenergic receptor gene under conditions allowing the expression of such gene is performed. The expression product, which may be exposed at the surface of the transformed cell host or present in the cytoplasm of the host cell, is then utilized to test a variety of ligands, drugs and chemical agents by placing these agents in contact with the transformed cell host. An affinity reaction between the transformed cell host and the specific agent is then detected.

[0052] The transformed host cells containing the polypeptide sequence of the β 3-adrenergic receptor can also be used to test intracellular responses to various chemical agents.

[0053] More specifically, it is known in the art that a variety of hormones, both peptidic and non-peptidic, exert their cellular effects through interaction with cell surface receptors

coupled to guanine nucleotide binding proteins or the G-proteins. The G-protein coupled receptors mediate a variety of intracellular responses, wherein the initial specificity is usually determined at the level of the cell surface receptor. The diversity of the cellular responses results from the differential coupling of the receptor to various G-proteins, each of which stimulates a distinct intracellular effector system. The G-protein linked receptor can bind a variety of agonists which stimulates the G protein coupled to the receptor. The activation of the stimulatory G protein leads in turn to adenylate cyclase stimulation and thus an increase in cAMP levels in cells. The β 3-adrenergic receptor has the ability to couple G protein and thus the effects of various agonists to activate the stimulatory G protein, raising cAMP levels in the cell can also be tested using the transformed cell hosts of the present invention.

[0054] It is of extreme importance that the entire gene sequence be utilized to conduct the above-identified studies, since it is known that the additional sequences described in the present invention effect the regulation of these receptors and especially influence the receptor's expression. Thus, for instance, to test specific ligands and drugs which may aid in an agent to reduce obesity (since the β 3-adrenergic receptor is present in low quantities in obese patients) it may be necessary to increase the amount of β 3-receptor present. The RNA expression levels of the β 3-adrenergic receptor may be regulated by additional sequences described and characterized in the present invention such as the sequences present in the promoter region, the intron/exon region and the 3' untranslated region. Thus, an agent that can be found to increase the transcription of β 3-receptors may lead to treatment for obesity and diabetes.

[0055] The cloning and initial sequencing of the human and mouse β 3-adrenergic receptor genomic genes have been previously described. Nahmias et al., THE EMBO JOURNAL vol. 16, No. 12 pgs. 3721-3727 (1991) and Emorine et al., Science vol. 245, pgs. 1118-1121 (1989). Both of these references are incorporated herein by reference. To obtain more insight into the mechanisms at the basis of the regulated expression of the β 3-adrenergic receptor in both human and mouse, further study was undertaken. By comparing the β 3-adrenergic receptor gene and the PCR-amplified cDNA sequences, the presence of introns which interrupted the coding and the 3'-untranslated regions of the human and mouse β 3-adrenergic receptor gene was discovered.

[0056] Nucleotide sequencing was performed on a 2.5 kb SphI-BglII fragment spanning the entire 3'-untranslated regions of the human gene and a 2.2 kb XhoI-BglII fragment spanning the entire 3' untranslated region of the mouse β 3-genomic genes after insertion of these fragments in both orientations into M13 cloning vectors. Individual subclones were then sequenced by the standard chain termination method. The primers used were oligonucleotides synthesized step by step during the course of the sequence determination.

[0057] A preferred embodiment of the present invention is the identification of two exons in humans and three exons in mice, which contain the entire β 3-adrenergic receptor gene sequence. In humans, a large 1.4 kb exon encodes the first 402 amino acid residues, while the second exon of 700 base pairs encodes the sequence for the six carboxy-terminal residues of β 3-adrenergic receptor, as well as the entire untranslated 3' region of the β 3-adrenergic receptor mRNA.

[0058] In mice, there are three exons. The first exon encodes 388 amino acid residues of the β 3-adrenergic receptor; a second exon of 68 base pairs encodes for the twelve carboxy-terminal residues of the β 3-adrenergic receptor and the third exon of about 600 to 700 base pairs with two acceptor splice sites codes for the 3'-untranslated region of β 3-adrenergic receptor mRNA. In mouse adipose tissues, alternate splicing at the two splice sites found in the third exon generates two β 3-adrenergic transcripts. The shorter transcript is the major transcript found in white adipose tissue, while the longer transcript is slightly predominant in brown adipose tissue.

[0059] Thus, the amino acid sequence for human β 3-adrenergic receptor is 6 amino acids longer than previously thought, while the mouse sequence is 12 amino acids longer than previously reported. The nucleotide and amino acid sequences as well as the mRNA splicing are depicted in FIGS. 1-5 for the human and mouse β 3-adrenergic receptor.

[0060] The localization of the mRNA transcription start sites was undertaken by S1 nuclease mapping and the mRNA transcription start sites for both mouse and human β 3-adrenergic receptor mRNA were then localized in a region between 150-190 nucleotides upstream from the ATG translation start cocon. In mouse, a second mRNA initiation region exists further upstream. Further upstream from the mRNA cap sites, glucocorticoid response

elements were found in the immediate vicinity of recognition sequences for transcription factor AP-1. The inhibitory effects of glucocorticoids on β 3-adrenergic receptor mRNA levels may result from interactions of glucocorticoid receptor with c-fos and c-jun products which constitutes transcription factor AP-1.

[0061] Thus, the entire genes including the promoter region, the intron/exon region and the 3'-untranslated region, coding for β 3-adrenergic receptors were obtained, as well as the mRNA transcription start sites. This finding is of utmost importance not only for detection and diagnostic purposes, but also to obtain drugs useful in the regulation of β 3 expression in tissue. Thus, the utilization of alternate promoters and/or splice sites allows tissue specific regulation of β 3-adrenergic expression.

EXAMPLES

EXAMPLE 1

Isolation and sequencing of the murine β 3AR gene

[0062] All the methods used for recombinant DNA procedures are from Ausubel et al. Science 245, pgs 1118-1121 (1987). A 1.3 kb NcoI-BamHI DNA fragment encompassing the entire coding region of the human β 3AR gene (Emorine et al. Current Protocols in Molecular Biology, Greene publishing Associates and Wiley InterScience, New York (1989)) was used as a probe to screen a mouse NIH 3T3 genomic library in the lambda FIX.TM.II vector (Stratagene). The nitrocellulose replica filters of the library were prehybridized for 16 h at 42° C. in a buffer containing 8 mM Tris-HCl (pH 7.5), 40% formamide, 4.times.SSC, 5.times.Denhardt's, 0.2% SDS, 50 mM sodium phosphate and 100 .mu.g/ml heat-denatured salmon sperm DNA. Hybridization was performed for 16 h at 42° C. in the same buffer containing the probe (2×10^6 c.p.m./ml) 32 P-labelled by the random priming method to a specific activity of 10^9 c.p.m./ μ g. Final washes were at 45° C. in 0.1 x SSC and 0.05% SDS.

[0063] A 2 kb BglII-BamHI restriction fragment hybridizing to the probe was subcloned in both orientation into M13mp18. Nested deletions were created with the exonuclease III and sequencing was performed by the dideoxy chain termination method using [α - 35 S]dATP

(800 Ci/mmol, Amersham) and *Thermophilus aquaticus* (Taq) polymerase (Taquence kit, USB). Sequence analysis were carried out using the CIT12 (University Paris V, Paris, France) computer software facilities.

EXAMPLE 2

DNA probe and Hybridization Analysis

[0064] A 310 bp DNA fragment ("A43") corresponding to the N-terminus of the mouse β 3-adrenergic receptor was produced by polymerase chain reaction using a template of 5 ng of the lambda clone containing the mouse β 3-adrenergic receptor gene. Thirty cycles of amplification at 93° C. for 1.5 minute; 55° C. for 2 minutes; and 72° C. for 2 minutes were performed using 2.5 U Taq polymerase (Cetus) in 100 μ l of buffer containing 10 mM Tris-HCl (pH 8.4), 3 mM MgCl₂, 0.05% TWEEN 20, 0.05% NP40, 10% dimethylsulphoxide, 5% formamide, 125 μ M of each deoxynucleotide triphosphate and 125 pM of each primer. Sense (SEQ ID NO:7) (5'-GCTCCGTGGCCTCACGAGAA-3') and antisense (SEQ ID NO:8) (5'-CCCAACGGCCAGTGGCCAGTCAGCG-3') primers corresponding to amino acids 2-8 and 98-106, respectively of the human β 3-adrenergic receptor sequence. The amplified 310 bp DNA fragment was purified by electrophoresis through an acrylamide gel. ³²P-labelled by random priming to a specific activity of 10⁹ c.p.m./ μ g, and used as a probe in Southern and Northern blot experiments.

[0065] For Southern blot hybridization analysis, single restriction enzyme digests of genomic DNA (10 μ g) were performed and the samples were electrophoresed through a 0.8% agarose gel and blotted onto nylon (Hybond N+, Amersham) filters. Prehybridization and hybridization procedure were performed as described above and final washes were at 45° C. in 0.1 x SSC, 0.05% SDS.

[0066] For Northern blot analysis, total RNA extracted from mouse or cultured cells was denatured in Glyoxal-DMSO, electrophoresed through a 1% agarose gel and transferred onto a nylon membrane (Hybond N+, Amersham). Prehybridization and hybridization, as well as washing were performed as described above, except that hybridization was performed in the presence of 10% dextran sulphate.

EXAMPLE 3

In situ hybridization

[0067] For chromosomal localization of the β 3-adrenergic receptor gene in mouse, in situ hybridization was performed on metaphase spreads of concavalin A-stimulated lymphocytes from a WMP male mouse, in which all the autosomes except number 19 were in the form of metacentric Robertsonian translocations. For localization of the β 3-adrenergic receptor gene in man, metaphase spreads were from phytohaemagglutinin-stimulated human lymphocytes. In both cases the lymphocytes were cultured at 37° C. for 72 hrs., with 5-bromodeoxyuridine (60 μ g/ml) added for the final 7 hours of culture.

[0068] Specific probes were the 310 bp A43 fragment of the mouse β 3-adrenergic receptor gene and a 206 bp (AccI-ApaLI) DNA fragment encompassing the third intracytoplasmic loop of the human β 3-adrenergic receptor. These fragments were subcloned into the pUC19 plasmid vector, tritium labelled by nick translation to a specific activity of 10^8 d.p.m./ μ g, and used at a final concentration of 25 ng/ml of hybridization solution. Hybridization to metaphase spreads, post-hybridization washes, emulsion autoradiography, R-banding and silver grain analysis were carried out as described by Mattei et al. HUMAN GENETICS, 69, pgs. 268-271 (1985).

EXAMPLE 4

Construction, cell culture and transfections

[0069] For expression in eukaryotic cells, a 1365 bp NarI-BamHI restriction fragment from the mouse β 3-adrenergic receptor gene was inserted under the control of the SV40 early promoter into a plasmid vector containing the 3'-untranslated region of the gene for the hepatitis-B surface antigen, and murine dihydrofolate reductase (DHFR) gene as a selectable marker as described by Larsky et al. Biotechnology, 2, pgs. 527-530 (1984). The resulting construct contained 15 bp of 5' untranslated region, 1164 bp of coding region and 186 bp of 3' non-coding region of the mouse β 3-adrenergic receptor gene.

[0070] CHO cells, deficient in DHFR, were grown in Ham's F12 medium (Seromed) supplemented with 10% fetal calf serum, 2 mM L-glutamine, 15 µg/ml glycine, 9.5 µg/ml hypoxanthine and 1.46 µg/ml thymidine. Transfection with 5 µg of the expression vector was performed by the calcium phosphate precipitation method as described by Tate et al. EUR. J. Biochem., 196, pgs. 357-361 (1991). Stable transformants expressing the DHFR gene were screened for the expression of β adrenergic receptor by stimulation of cAMP accumulation with 1 µM (-)-isoproterenol. One clone giving the highest stimulation was subcloned by limiting dilution, and is referred to as CHO-Mo β 3'.

EXAMPLE 5

Isolating and sequencing of the human β 3AR gene.

[0071] Similar procedures set forth in the above examples to isolate and sequences the murine β 3-adrenergic receptor gene were used in the isolation of the human β 3-adrenergic receptor gene. A human genomic library was screened with the entire coding regions of the gene for the β 1-adrenergic receptor of turkey Yarden et al., P.N.A.S., 83, 6795 (1986) and of the human gene for the β 2-adrenergic receptor Emorine et al P.N.A.S. 84, 6995 (1987). The library was constructed in the BamHI sites of the EMBL4 vector from size-selected (15 to 20 kb) fragments of human placental DNA partially digested with Sau3a. For plaque purification, probes of a 1.3 kb NcoI-AhaI restriction fragment of the human β 2-adrenergic receptor and the 1.8 kb fragment of the turkey β 1-adrenergic receptor were used Yarden et al., P.N.A.S., 83, 6795 (1986).

[0072] Among 43 positive clones two carried the gene coding for the human β 1-adrenergic receptor and another two the gene for the β 2-adrenergic receptor.

[0073] A family of 14 homologous clones consisting of a group of six identical clones, a second group of three identical clones and a third group of two identical clones and three independent clones displayed sequences homologous to both probes in a 2.1 kb BamHI and BglII fragment. From one clone, this 2.1 kb fragment was sequenced using the methods described by the chain termination method of Sanger et al., P.N.A.S., 74, 5463 (1977). This

fragment that was entirely sequenced and shown to contain a gene coding for a polypeptide of 402 amino acids with a predicted size of 42,881 daltons.

EXAMPLE 6

Nucleotide sequence determination

[0074] 2.5 kb SphI-BglII, and 2.2 kb XhoI-BglII fragments spanning the entire 3'-untranslated regions of the human and mouse β 3-AR genomic genes, respectively, were inserted in both orientations into M13 and individual subclones were sequenced by the standard termination method (Taquence DNA sequencing kit, USB, Ohio). Primers were oligonucleotides synthesized step by step during the course of sequence determination.

EXAMPLE 7

PCR-generated cDNA

[0075] Total RNA was prepared from frozen-powdered tissues and digested for 1 hr at 37° C. with 0.3 U of RNase-free DNase I (Promega, Wisconsin) per μ g of nucleic acid in 100 mM Tris-HCl, pH 7.5; 50 mM MgCl₂, in the presence of 2 U/ml of placenta RNase inhibitor. 0.25 μ g of RNA was then treated with 400 U of maloney murine leukemia virus reverse transcriptase (Gibco BRL) in 20 μ l of PCR buffer containing 67 mM Tris-HCl, pH 8.4; 6.7 mM MgCl₂ ; 6.7 mM EDTA; 10 mM β -2-mercaptoethanol; 16 mM (NH₄)₂SO₄ ; and 0.1 mg/ml gelatine containing 0.4 mM each of dNTP, 10 mM random hexanucleotides and 2 U/ml RNase inhibitor. A control without reverse transcriptase was also run to ensure that amplification of the nucleotide sequence did not proceed from residual genomic DNA. 80 μ l of PCR buffer containing 2.5 U of Thermophilus aquaticus polymerase (Perkin-Elmer-Cetus), 125 nM each of sense and antisense oligonucleotide primers, containing the following sequences, respectively:

[0076] 5' GCTCCGTGGCCTCACGAGAA 3' (SEQ ID NO:7) and

[0077] 5' CCCAACGGCCAGTGGCCAGTCAGCG 3' (SEQ ID NO:8),

[0078] 125 mM each dNTP and 10% (v/v) dimethylsulfoxide was added to the cDNA samples which were submitted to 29 cycles at temperatures of 92° C., 1 minute; 57° C., 1.5 minutes; 72° C.; 1.5 minutes; followed by 7 minutes of extension at 72° C. in a temperature cyclers (LEP-PREM).

EXAMPLE 8

Determination of mRNA transcription start sites

[0079] For nuclease S1 mapping single stranded DNA from a M13 subclone of the murine β 3-AR gene promoter region (fragment BglIII-NarI, was used as template for probe synthesis. After polymerization in the presence of α [³²P]-dATP, DNA was digested with Alw NI and the single stranded probe (380 nucleotides including 310 base from the β 3-AR gene plus 70 bases from M13) was isolated on denaturing 6% polyacrylamide gels containing 7 M urea. Probe (10^5 c.p.m.) and total RNA (50 μ g) were mixed in 30 μ l of 40 mM Pipes buffer pH 6.4, containing 0.4 M NaCl, 1 mM EDTA and 70% formamide and denatured for 10 min at 85° C. After hybridization (14-16 hr at 30° C.), samples were digested for 30 min at 37° C. with 100-200 units of S1 nuclease in 400 μ l of 50 mM sodium acetate, pH 4.8, 280 mM NaCl, 4.5 mM ZnSO₄. Following incubation for 10 min at 65° C., samples were phenol-extracted, concentrated by ethanol precipitation and analyzed on 6% acrylamide, 7 M urea sequencing gels.

[0080] For primer extension, oligonucleotide A74 (36-mer) having the following sequence (SEQ ID NO:9):

[0081] 5' CTGGTCCAGGGGAGGGGACAGCAAGGCATGAGAGCG 3'

[0082] was labeled with T4-polynucleotide kinase hybridized (5×10^5 c.p.m.) to 50 μ g of total RNA, in 30 μ l of 20 mM Tris-HCl, pH 8, 50 mM NaCl, 10 mM MgCl₂, 1 mM dithiothreitol, and 0.1 mM EDTA, for 14-16 hr at 37° C. Nucleic acids were ethanol precipitated and incubated for 90 min at 42° C. after resuspension in 25 μ l of 50 mM Tris-HCl, pH 8.3, 8 mM MgCl₂, 30 mM KCl containing 0.5 mM each DNTP, 50 units of placenta

09395241-103404
T01E01-T25660

RNase inhibitor, and 40 units of avian myeloblastosis virus reverse transcriptase. Reactions were terminated by addition of 1 μ l of both 0.5 M EDTA and pancreatic RNase A (1 mg/ml) and incubation for 30 min at 37° C. was performed. After phenol-extraction and ethanol precipitation in the presence of ammonium acetate (2 M final) samples were analyzed on 6% acrylamide, 7 M urea sequencing gels.

EXAMPLE 9

Expression of the entire human β 3-adrenergic receptor

[0083] For expression in eukaryotic cells, a 3.7 kb SmaI-BglII restriction fragment from the human β 3-adrenergic receptor gene was inserted under the control of the SV40 early promoter into a plasmid vector also containing the murine dihydrofolate reductase (DHFR) gene as a selectable marker as described by Larsky et al. Biotechnology, 2, pgs. 527-530 (1984). The resulting construct contained 4 bp of 5' untranslated region, the entire intron/exon sequence of the human β 3-receptor gene with its own 3' untranslated region.

[0084] CHO cells or L cells (wherein the splicing of the gene can take place) deficient in DHFR, were grown in Ham's F12 medium (Seromed) supplemented with 10% fetal calf serum, 2 mM L-glutamine, 15 μ g/ml glycine, 9.5 μ g/ml hypoxanthine and 1.46 μ g/ml thymidine. Transfection with 5 μ g of the expression vector was performed by the calcium phosphate precipitation method as described by Tate et al. EUR. J. Biochem., 196, pgs. 357-361 (1991). Stable transformants expressing the DHFR gene were screened for the expression of β adrenergic receptor by stimulation of cAMP accumulation with 1 μ M (-)-isoproterenol.

EXAMPLE 10

Testing of various chemicals for β 3-adrenergic receptor affinity

[0085] Preconfluent cells (5×10^5) were incubated at 37° C. for 20 minutes in 0.5 ml of Hank's buffer containing 20 mM HEPES (pH 7.4), 1 mM ascorbic acid, 1 mM isobutylmethylxanthine and various concentrations of agonists such as BRL 37344, (-)-isoproterenol and (-)-norepinephrine. After boiling for 5 minutes and centrifuging at 4,000

r.p.m. for 10 minutes at 4° C., the amount of cAMP produced was measured using the Amersham cAMP assay kit. Inhibition studies for cAMP accumulation were carried out by preincubating cells at 37° C. for 10 minutes with an antagonist such as propranolol, ICI 118551 and CGP 20712A before the addition of 5 nm (-)-isoproterenol. The mixture was incubated for an additional 20 minutes. Computer analysis of the data was carried out using the Graph-PAD program (copyright 1987 by M. J. Motulsky).

[0086] The entire disclosure of each document cited (including patents, patent applications, journal articles, abstracts, or other disclosures) in the Background of the Invention, Detailed Description, and Examples is herein incorporated by reference. Further, the hard copy of the sequence listing submitted herewith and the corresponding computer readable form are both incorporated herein by reference in their entireties.

[0087] Certain β 3 Adrenergic Receptor Gene polynucleotides and polypeptides of the invention were disclosed in U.S. application 08/450,962, filed May 25, 1995, now U.S. Patent No. 6,274,706, which is herein incorporated by reference in its entirety. Moreover, the hard copy and computer readable form of the Sequence Listing of U.S. application 08/450,962, filed May 25, 1995, now U.S. Patent No. 6,274,706, are herein incorporated by reference in their entireties.